A Guide to First-Line Protocol to Common Urinary/Men's Sexual Health Conditions



ERECTILE DYSFUNCTION

First-line protocol:

- Check testosterone level
- Trial Sildenafil and/or Tadalafil

Advanced treatments from Spokane Urology:

- Hormone therapy
- Penile implant
- Vascular surgery

INCONTINENCE

Symptoms: leakage from stress event (sneeze, cough, exercise) or urge (immediate need to go)

First-line protocol:

- Pelvic floor physical therapy or Kegel exercises
- Dietary changes to reduce irritants (caffeine, alcohol, spicy)
- Trial Oxybutynin or other OAB med
- Pessary ring

Advanced treatments from Spokane Urology:

- BOTOX® injection
- Electronic stimulation (InterStim™)
- Transvaginal sling
- Urethral bulking agent (Bulkamid®)

INTERSTITIAL CYSTITIS

Symptoms: general symptoms like bladder pain, dysuria, pelvic pain

First-line protocol:

- Pelvic floor physical therapy or Kegel exercises
- Dietary changes to reduce irritants (caffeine, alcohol, spicy)
- OTC anti-inflammatories

Advanced treatments from Spokane Urology:

- Cystoscopic to evaluate the bladder and ensure there are no causes
- Bladder hydrodistention to diagnose and treat through stretching
- BOTOX® injection

OVERACTIVE BLADDER

Symptoms: urinary frequency or urgency

First-line protocol:

- Pelvic floor physical therapy or Kegel exercises
- Dietary changes to reduce irritants (caffeine, alcohol, spicy)
- Ensure good bowel regimen to remove constipation as a trigger
- Trial Oxybutynin or other OAB med

Advanced treatments from Spokane Urology:

- BOTOX® injection
- Electronic stimulation (InterStim™)
- Transvaginal sling
- Urethral bulking agent (Bulkamid®)
- Percutaneous tibial nerve stimulation (PTNS)

TESTICULAR PAIN

First-line protocol:

- Supportive underwear/athletic supporter
- Ice pack
- Trial nonsteroidal anti-inflammatory drugs (NSAID)
- Send for testicular ultrasound prior to referral

Advanced treatments from Spokane Urology:

• Evaluation for underlying diseases or testicular cancer

URINARY TRACT INFECTION

Symptoms: recurring; more than 3 in 12 months

First-line protocol:

- Hygiene discussion
- Increase hydration (minimize caffeine and alcohol)
- Evaluate for bowel control issues of constipation or diarrhea
- Trial of cranberry supplements, d-mannose, and vaginal estrogen
- Short-term course of antibiotics

Advanced treatments from Spokane Urology:

- Longer-term course of antibiotics
- Evaluation for underlying diseases or medications compromising the immune system

NEPHROLITHIASIS (KIDNEY STONES)

Symptoms: flank pain; cloudy urine; urinary frequency; history of stones **First-line protocol**:

• Send for imaging to confirm presence of stones prior to referral

Advanced treatments from Spokane Urology:

- Lithotripsv
- Cysto, ureteroscopy, laser litho, retrograde pyelogram, stent placement
- Cysto, ureteroscopy, laser litho, stone basketing, stent placement
- Cystolitholapaxy for bladder stone
- Extracorporeal shockwave lithotripsy (ESWL)
- Percutaneous nephrolithotomy (PCNL)

The providers at Spokane Urology are here to help your patients who have persistent conditions that may require advanced therapies after trying first-line protocols.

If you have questions about when to refer a patient or any of the advanced treatments we provide, please contact:

Kristin Brown, Chief Operating Officer, Spokane Urology brownk@spokaneurology.com | (509) 747-3147 EXT 132

MEET OUR UROLOGISTS



Evan Carlos, MD



Levi Deters, MD, FACS



Raymond Lance, IVIL



Michael Maccini, MD




Bradford Nelson, MD, PharmD



Shane Pearce, MD



Duncan Sloan, MD



Bryan Voelzke, MD, FACS

PHYSICIAN OFFICES

1401 East Trent Ave. Suite 200 Spokane, WA 99202 12525 E. Mission Ave. Suite 202 Spokane Valley, WA 99216

OUTPATIENT SURGERY CENTER

1401 East Trent Ave. Suite 100 Spokane, WA 99202